
The effect of cold atmospheric pressure plasma (CAP) on cell migratory behaviors and molecular markers of wound healing machinery

Debarati Shome¹, Niklas Hohmann¹, J Schmidt², Thomas von Woedtke¹, and Kai Masur¹

¹Leibniz Institute for Plasma Science & Technology, Germany

²Clinic Karlsburg, Dr. Guth GmbH & Co. KG, Karlsburg, Germany

Abstract:

Cold atmospheric pressure plasma (CAP) is a promising tool for biomedical and clinical application. Cold physical plasma are partially ionized gases that mediate biological response generating ROS and RNS species.¹⁻³ In this study, we used the medical device class 2a kINPen MED®; it is clinically approved atmospheric argon plasma jet. Typical active agents being generated include ions, electrons, and reactive oxygen and nitrogen species (ROS/RNS). Electric and magnetic fields, light (visible, in-fared, UV), and neutral particles are also being generated⁴. It is well known, that wound oxygenation is an important factor of wound healing and scavenging reactive species could impair wound healing.⁵ In this study, we examined the extent of wound healing and the underlying cellular mechanism in vitro induced by CAP. Our group had collected wound exudates before and after CAP treatment from 7 ambulant diabetic patients with chronic wounds from 'Competence Center Diabetes' in Karlsburg. These wound exudates were also analyzed.

Methods: For our studies, we incorporated dermal keratinocytes, fibroblasts and co culture. An indirect treatment where CAP treated media (RPMI) was added and a direct CAP treatment on the cells (different time points) were performed and the migration assay was monitored. Matrix metalloproteinase play a pivotal role in wound re epithelization. The amount of metalloproteinase in the patient exudates were also monitored by ELISA. The gene expression upon CAP treatment was monitored with qPCR.

Results and future direction: Short-term CAP treatment induces enhanced cell migration than longer treatment and compared to untreated control. Co culture studies show an improved cell migration upon CAP treatment compared to keratinocytes alone. Matrix metalloproteinases were also reduced in wound exudates after CAP treatment which indicates improved wound healing. It was also evident with an average of 70-80 % reduction in the wound size of the patients. The signalling machinery of wound healing involving inflammatory (interleukin mediated), regenerative pathways (mediated by HIPPO signalling) and growth factors (mediated by IGF) are currently being checked to identify an autocrine/paracrine signalling pathway induced by CAP.

References:

1. Jablonowski, H. and T. von Woedtke (2015). "Research on plasma medicine-relevant plasma-liquid interaction: What happened in the past five years?" Clinical Plasma Medicine 3(2): 42-52.

2. Reuter, S., et al. (2012). "From RONS to ROS: Tailoring Plasma Jet Treatment of Skin Cells." IEEE Transactions on Plasma Science 40(11): 2986-2993.

3 David, B. G. (2012). "The emerging role of reactive oxygen and nitrogen species in redox biology and some

implications for plasma applications to medicine and biology." Journal of Physics D: Applied Physics 45(26): 263001.

4. Bekeschus, S., et al. (2016). The plasma jet kINPen – A powerful tool for wound healing. Clinical Plasma Medicine . 4: 19-28.

5. Schmidt, A., et al. (2017). "One Year Follow-Up Risk Assessment in SKH-1 Mice and Wounds Treated with an Argon Plasma Jet." International Journal of Molecular Sciences 18(4):868.